

AMENDMENTS

SPECIFICATION

Please replace the paragraph beginning on page 19, line 11 and ending on page 20, line 7 (as previously requested in Applicants' Response dated September 8, 1998) with the following paragraph:

Compounds particularly useful according to the present invention are extendin agonist compounds of the formula (I):

1 5 10

Xaa₁ Xaa₂ Xaa₃ Gly Thr Xaa₄ Xaa₅ Xaa₆ Xaa₇ Xaa₈

15 20

Ser Lys Gln Xaa₉ Glu Glu Glu Ala Val Arg Leu

25 30

Xaa₁₀ Xaa₁₁ Xaa₁₂ Xaa₁₃ Leu Lys Asn Gly Gly Xaa₁₄

35

Ser Ser Gly Ala Xaa₁₅ Xaa₁₆ Xaa₁₇ Xaa₁₈ -Z

wherein Xaa₁ is His, Arg or Tyr; Xaa₂ is Ser, Gly, Ala or Thr; Xaa₃ is Asp or Glu; Xaa₄ is Phe, Tyr or naphthylalanine; Xaa₅ is Thr or Ser; Xaa₆ is Ser or Thr; Xaa₇ is Asp or Glu; Xaa₈ is Leu, Ile, Val, pentylglycine or Met; Xaa₉ is Leu, Ile, pentylglycine, Val or Met; Xaa₁₀ is Phe, Tyr or naphthylalanine; Xaa₁₁ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met; Xaa₁₂ is Glu or Asp; Xaa₁₃ is Trp, Phe, Tyr, or naphthylalanine; Xaa₁₄, Xaa₁₅, Xaa₁₆ and Xaa₁₇ are independently Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine or N-alkylalanine; Xaa₁₈ is Ser, Thr or Tyr; and Z is -OH or -NH₂; with the proviso that the compound does not have the formula of either SEQ. ID. NOS. 1 or 2. Preferred N-alkyl groups for N-alkylglycine, N-alkylpentylglycine and N-alkylalanine include lower alkyl groups preferably of 1 to about 6 carbon atoms, more preferably of 1 to 4 carbon atoms. Suitable compounds include those having amino acid sequences of SEQ. ID. NOS. 5 to 35.